

Les mardis de la grenouille

January 2024 - Webinar

FROGSSTAT Tools

LUCAS AUER, MARIA BERNARD, LAURENT CAUQUIL, MAHENDRA MARIADASSOU, GÉRALDINE PASCAL

How different
are two
communities?

My samples
are they
homogenous
or diverse?

What is the
composition of
each
community?

Are the communities
structured by some known
environmental factor (pH,
height, etc)?

Are there
interactions
between
species and
communities?

Are there ASV with
differential
abundance between
conditions?

FROGSSTAT with Phyloseq R package

- R package (McMurdie and Holmes, 2013) to analyse community composition data in a phylogenetic framework

It uses other R packages:

- Community ecology functions from vegan, ade4
- Tree manipulation from ape
- Graphics from ggplot2
- Differential analysis from DESeq2

Exercise 1

→ At the end of FROGS pipeline, what kind of data do we have ?

Exercise 1

→ At the end of FROGS pipeline, what kind of data do we have ?

FROGS biom containing:

- ASV count tables (required)
- ASV description : taxonomy

Phylogenetic tree in Newick format

Metadata: sample description in TSV file

Exercise 1

→ Take a look at the metadata

Exercise 1

→ Take a look at the metadata

FoodType:

Meat or Seafood

EnvType: 8 environment types

Meat → Ground Beef, Ground veal, Poultry sausage, Diced bacon

Seafood → Cooked schrimps, Smoked salmon, Salmon filet, Cod filet

	EnvType	Description	FoodType
	EnvType	Description	FoodType
DLT0.LOT01	DesLardons	LOT1	Meat
DLT0.LOT03	DesLardons	LOT3	Meat
DLT0.LOT04	DesLardons	LOT4	Meat
DLT0.LOT05	DesLardons	LOT5	Meat
DLT0.LOT06	DesLardons	LOT6	Meat
DLT0.LOT07	DesLardons	LOT7	Meat
DLT0.LOT08	DesLardons	LOT8	Meat
DLT0.LOT10	DesLardons	LOT10	Meat
MVT0.LOT01	MerguezVolaille	LOT1	Meat
MVT0.LOT03	MerguezVolaille	LOT3	Meat
MVT0.LOT05	MerguezVolaille	LOT5	Meat
MVT0.LOT06	MerguezVolaille	LOT6	Meat
MVT0.LOT07	MerguezVolaille	LOT7	Meat
MVT0.LOT08	MerguezVolaille	LOT8	Meat
MVT0.LOT09	MerguezVolaille	LOT9	Meat
MVT0.LOT10	MerguezVolaille	LOT10	Meat
BHT0.LOT01	BoeufHache	LOT1	Meat
BHT0.LOT03	BoeufHache	LOT3	Meat

Phyloseq Import Data tool

PHYLOSEQ OBJECT CREATION

Phyloseq : Data import

1. Statistical analysis is done in R, so an R object called Rdata must be created.

2. Run PhyloSeq Data import

The FROGS biom format contains:

- ASV count tables (required)
- ASV description : taxonomy

Others information used in FROGSSTAT are:

- sample description in TSV file
- phylogenetic tree in Newick format (nwk or nhx)

3. Create 2 phyloseq objects, with and without normalization (rename them)

FROGSSTAT Phyloseq Import Data from 3 files: biomfile, samplefile, treefile
(Galaxy Version 4.1.0+galaxy1) ☆ Favorite 🔄 Versions ▼ Options

Abundance biom file with taxonomical metadata (format: BIOM)

1: FROGS_5 Taxonomic affiliation: affiliation_abundance.biom

The file contains the ASV information (--biomfile)

Metadata associated to samples (format: TSV)

3: metadata_chaillou.tsv

The file contains the metadata that characterise each sample. (--samplefile)

Taxonomic tree file (format: Newick)

2: FROGS Tree: tree.nwk

The file contains the taxonomic tree information from FROGS Tree tool (optional) (--treefile)

Names of taxonomic levels

Kingdom Phylum Class Order Family Genus Species

The ordered taxonomic levels stored in BIOM. Each level is separated by one space (--ranks)

Do you want to normalise your data ?

No, keep abundance as it is.
 Yes, subsample abundances to the smallest sample size.

To normalise data before statistical analysis (default : No) (--normalisation)

Email notification

No

Send an email notification when the job completes.

Exercise 2

1. What are the resulting datasets ?
2. What is the difference between the resulting objects with and without normalization ?
3. Explore the HTML results

Exercise 2

1. What are the resulting datasets ?

- asv_data.Rdata file: R object used by phyloseq package for statistics
- HTML report: summary of the phyloseq object

Exercise 2

2. What is the difference between the resulting objects with and without normalization ?

Without normalization



Summary

Ranks Names

Sample metadata

Plot tree

Code

```
phyloseq-class experiment-level object
otu_table()  OTU Table:          [ 495 taxa and 64 samples ]
sample_data() Sample Data:       [ 64 samples by 4 sample variables ]
tax_table()  Taxonomy Table:    [ 495 taxa by 7 taxonomic ranks ]
phy_tree()   Phylogenetic Tree: [ 495 tips and 494 internal nodes ]
```



ASV are still called OTU in phyloseq functions

Exercise 2

2. What is the difference between the resulting objects with and without normalization ?

Summary

Ranks Names

Sample metadata

Plot tree

Code

```
phyloseq-class experiment-level object
otu_table() OTU Table: [ 495 taxa and 64 samples ]
sample_data() Sample Data: [ 64 samples by 4 sample variables ]
tax_table() Taxonomy Table: [ 495 taxa by 7 taxonomic ranks ]
phy_tree() Phylogenetic Tree: [ 495 tips and 494 internal nodes ]
```

Code

```
Number of sequences in each sample after normalization: 7638
```

With normalization (rarefaction)

Minimum number of sequences kept in each sample



Exercise 2

2. What is the difference between the resulting objects with and without normalization ?

With normalization (rarefaction)



Be aware that the number of taxa may decrease due to normalization

Summary

Ranks Names

Sample metadata

Plot tree

Code

```
phyloseq-class experiment-level object
otu_table() OTU Table: [ 495 taxa and 64 samples ]
sample_data() Sample Data: [ 64 samples by 4 sample variables ]
tax_table() Taxonomy Table: [ 495 taxa by 7 taxonomic ranks ]
phy_tree() Phylogenetic Tree: [ 495 tips and 494 internal nodes ]
```

Code

```
Number of sequences in each sample after normalization: 7638
```

Exercise 2

3. Explore the HTML results

Phyloseq 1.20.0



Code

Summary

Ranks Names

Sample metadata

Plot tree

Code

Taxonomic levels

```
Rank names : Kingdom, Phylum, Class, Order, Family, Genus, Species
```

Exercise 2

3. Explore the HTML results

Summary Ranks Names **Sample metadata** Plot tree

Sample variables: EnvType, Description, FoodType, SampleID

Code

EnvType : DesLardons, MerguezVolaille, BoeufHache, VeauHache, SaumonFume, FiletSaumon, FiletCabillaud, Crevette

Description : LOT1, LOT3, LOT4, LOT5, LOT6, LOT7, LOT8, LOT10, LOT9, LOT2

Code

FoodType : Meat, Seafood

SampleID : DLT0.LOT01, DLT0.LOT03, DLT0.LOT04, DLT0.LOT05, DLT0.LOT06, DLT0.LOT07, DLT0.LOT08, DLT0.LOT10, MVT0.LOT01, MVT0.LOT03, MVT0.LOT05, MVT0.LOT06, MVT0.LOT07, MVT0.LOT08, MVT0.LOT09, MVT0.LOT10, BHT0.LOT01, BHT0.LOT03, BHT0.LOT04, BHT0.LOT05, BHT0.LOT06, BHT0.LOT07, BHT0.LOT08, BHT0.LOT10, VHT0.LOT01, VHT0.LOT02, VHT0.LOT03, VHT0.LOT04, VHT0.LOT06, VHT0.LOT07, VHT0.LOT08, VHT0.LOT10, SFT0.LOT01, SFT0.LOT02, SFT0.LOT03, SFT0.LO

Variable names

Script R

the different modalities for each qualitative variable

Warning !
Metadata order (in each sample variable) are used to organize graphics.
So take extra care when you construct your sample_metadata file

It may make sense to order the metadata file i.e. the meats are together and the seafood together

Exercise 2

3. Explore the HTML results

Summary


Ranks Names

Sample metadata

Plot tree



Phylogenetic tree colored by Phylum



Phylum

- Actinobacteriota
- Bacteroidota
- Campylobacterota
- Cyanobacteria
- Desulfobacterota
- Firmicutes
- Fusobacteriota
- Patescibacteria
- Proteobacteria
- Spirochaetota

Exercise 2

3. Explore the HTML results

Summary

Ranks Names


Sample metadata

Plot tree

→ Information: Most represented phylum (in ASVs count)

- Bacteroidota
- Firmicutes
- Actinobacteriota
- Proteobacteria

Phylogenetic tree colored by Phylum



Biodiversity analysis

The points we will cover on biodiversity analysis

1. Exploring sample composition
2. Notions of biodiversity
3. α -diversity analysis
4. β -diversity analysis

I. Biodiversity analysis

COMPOSITION VISUALIZATION

Exploring biodiversity : visualization

FROGSSTAT Phyloseq Composition Visualisation with bar plot and composition plot (Galaxy Version 4.1.0+galaxy1)

Favorite

Versions

Options

Phyloseq object (format rdata)

4: FROGSSTAT Phyloseq Import Data SUBSAMPLED: asv_data.Rdata

This is the result of FROGS Phyloseq Import Data tool.

Grouping variable

EnvType

Experimental variable used to group samples (Treatment, Host type, etc). (--varExp)

Taxonomic level to filter your data

Kingdom

Ex: Kingdom, Phylum, Class, Order, Family, Genus, Species (--taxaRank1)

Taxa (at the above taxonomic level) to keep in the dataset

Bacteria

ex: Bacteria (when filtering at the Kingdom level), Firmicutes (when filtering at the Phylum level). More specific, i.e. Firmicutes Proteobacteria (--taxaSet1)

Taxonomic level used for aggregation

Phylum

Ex: Family (when filtering at the Phylum level). The aggregation level must be below the filtering level

Number of most abundant taxa to keep

9

Ex: 9, i.e. Tool keeps the 9 most abundant taxa and the remaining taxa are aggregated in a group 'Others'

Explore the sample **RAW** or **NORMALISED** count

Choose a sample variable to organize graphics: either EnvType or FoodType

At what taxonomic rank do we want to study?

Inside this taxonomic rank, what are the target group ?

On which rank do we want to group the ASVs?

Number of majority groupings to be displayed

For the first usage, let the default parameters

Exercise 3


1. What are the resulting datasets ?
2. What is the difference between Bar plot and Plot composition ?
3. What biological information could you extract ?
4. What are the perspectives for going further?

Exercise 3

1. What are the resulting datasets ?


→ HTML report: summary of the phyloseq object

- Bar plot
- Composition plot



Exercise 3


2. What is the difference between Bar plot and Plot composition ?



- one rectangle is one ASV
- one color is one phylum
- y axis: number of sequences – these are absolute counts
- size of rectangle depends on number of sequences

Exercise 3

2. What is the difference between Bar plot and Plot composition ?



Limitations:

- Plot bar works at the ASV-level and displays all the ASV at the specified rank
- This may lead to cluttered graphics and unnecessary legends
- No easy way to look at a subset of the data
- Works with absolute counts (beware of unequal depths or used normalized function)



[load-extra-functions.R](#)



Bar plot

Composition plot

Exploring biodiversity : visualization

Another graph: `plot_composition` function :

- Works with **relative abundances**
- Subsets ASVs** at a given taxonomic level
- Aggregates ASVs** at another taxonomic level
- Shows **only a given number** of taxa

Taxonomic level to filter your data

Kingdom

ex: Kingdom, Phylum, Class, Order, Family, Genus, Species

Taxa (at the above taxonomic level) to keep in the dataset

Bacteria

ex: Bacteria (when filtering at the Kingdom level), Firmicutes (when filtering at the Phylum level). Multiple taxa (separated by a space) can be specified, i.e. Firmicutes Proteobacteria

Taxonomic level used for aggregation

Phylum

ex: Family (when filtering at the Phylum level). The aggregation level must be below the filtering level.

Number of most abundant taxa to keep

9

ex: 9, i.e. Tool keeps the 9 most abundant taxa and the remaining taxa are aggregated in a group 'Other'


Exercise 3

Bar plot

Composition plot

2. What is the difference between Bar plot and Plot composition ?

- one rectangle is one phylum (no borderline) (or any other specified taxonomy rank)
- one color is one phylum
- y axis: counts are reduced to 1, so, here, we have relative counts




Exercise 3

Bar plot

Composition plot

3. What biological information could you extract ?

- Meat types on the left share common Phylum composition, with a majority of Firmicutes (easy to remark thanks of ordered levels)
- Seafoods seem to be much more variable
- Firmicutes and Proteobacteria are present in all samples, but with a wide range of abundance




Exercise 3

Bar plot

Composition plot

4. What are the perspectives for going further?




Exercise 3

Bar plot

Composition plot

4. What are the perspectives for going further?

- What is the composition of the 9 most abundant Families of *Firmicutes* ?
- What is the composition of the 9 most abundant Families of *Proteobacteria* ?



Exercise 4

1. What is the composition of the 9 most abundant Families of Firmicutes ?
2. What is the composition of the 9 most abundant Families of Proteobacteria ?

Exercise 4

1. What is the composition of the 9 most abundant Families of Firmicutes ?

Taxonomic level to filter your data

Phylum

ex: Kingdom, Phylum, Class, Order, Family, Genus, Species

Taxa (at the above taxonomic level) to keep in the dataset

Firmicutes

ex: Bacteria (when filtering at the Kingdom level), Firmicutes (when filtering at the Phylum level).

Multiple taxa (separated by a space) can be specified, i.e. Firmicutes Proteobacteria

Taxonomic level used for aggregation


Family

ex: Family (when filtering at the Phylum level). The aggregation level must be below the filtering level.

Number of most abundant taxa to keep

9


ex: 9, i.e. Tool keeps the 9 most abundant taxa and the remaining taxa are aggregated in a group 'Other'



Exercise 4

1. What is the composition of the 9 most abundant Families of Firmicutes ?

- Abundance does not reach 1 because only Phylum Firmicutes is displayed, the "missing" abundance is carried by other Phyla.
- As seen at the Phylum level, Firmicutes are more represented in meat types than in seafoods
- Dominant Firmicutes families are not the same in each food type



Exercise 4

2. What is the composition of the 9 most abundant Families of Proteobacteria ?

Taxonomic level to filter your data

Phylum

ex: Kingdom, Phylum, Class, Order, Family, Genus, Species

Taxa (at the above taxonomic level) to keep in the dataset

Proteobacteria

ex: Bacteria (when filtering at the Kingdom level), Firmicutes (when filtering at the Phylum level).

Multiple taxa (separated by a space) can be specified, i.e. Firmicutes Proteobacteria

Taxonomic level used for aggregation


Family

ex: Family (when filtering at the Phylum level). The aggregation level must be below the filtering level.

Number of most abundant taxa to keep

9

ex: 9, i.e. Tool keeps the 9 most abundant taxa and the remaining taxa are aggregated in a group 'Other'



Exercise 4


2. What is the composition of the 9 most abundant Families of Proteobacteria ?

- As seen at the Phylum level, Proteobacteria are particularly present in seafood samples
- SaumonFume samples with extremely high levels of Proteobacteria are dominated by Vibrionaceae family, while other food types are balanced between several families




Exploring biodiversity : visualization

Remark 1 : An example of what happens when sample metadata file is not sorted in a meaningful way



MEAT

SEAFOOD




disordered

Exploring biodiversity : visualisation

Remark 2 : Keep in mind that human eye cannot distinguish more than 12 colors at the same time.

Example of the 30 most abundant Families among Bacteria





II. Biodiversity analysis

DIVERSITY INDICES



Exploring biodiversity : descriptors

- The **richness** corresponds to the number of ASVs or functional groups present in communities. It characterizes the **composition**.
- The **diversity** takes into account the relative abundance of species. It characterizes the **structure**



Exploring biodiversity : descriptors

- The **richness** corresponds to the number of ASVs or functional groups present in communities. It characterizes the **composition**.
- The **diversity** takes into account the relative abundancy of species. It characterizes the **structure**




Richness : Eco1 = Eco2

Diversity: Eco2 > Eco1

Exploring biodiversity : statistical indices

3 levels of diversity:

- **α -diversity**: diversity **within** a community
- **β -diversity**: diversity **between** communities
 - β -dissimilarities/distances
 - dissimilarities between pairs of communities
 - often used as a first step to compute diversity
- γ -diversity: diversity at the landscape scale (blurry for bacterial communities)



Exploring biodiversity : statistical indices

There are qualitative, quantitative and phylogenetic indices:

Qualitative (Presence/Absence) vs. Quantitative (Abundance)

- Qualitative indices give equal weight to all species, dominant or rare
- Qualitative indices are more sensitive to differences in sampling depths
- Qualitative indices emphasize differences in taxa diversity while quantitative are more sensitive to increases in composition differences

Phylogenetic indices

- Require a phylogenetic tree
- phylogeny allows to attenuate clustering errors because 2 different ASVs can be phylogenetically close



III. Biodiversity analysis

α -DIVERSITY INDICES

4 α -diversity indices

1. Richness
2. Chao
3. Shannon
4. Inv-Simpson

Richness




Richness : Eco1 = Eco2

Richness
Number of observed species

α -diversity: Chao1

Richness	Chao
Number of observed species	Richness + (estimated) number of unobserved species



α -diversity: Chao1

Chao1 is an abundance-based estimator. This means that the data it needs relate to the abundance of taxa in the sample.

This index **estimates the number of unobserved species** from those that have only been **observed once or twice**. This diversity index is a minimum estimator. In order for it to fit the dataset, it is necessary that singletons and duplicates represent a significant part of the information

Many taxa, species, are represented by a few individuals (rare species) and others can be represented by many individuals (abundant species).

Well, **chao1 is based on the rare species**.

So we need to know how many species are represented by **1 individual (singleton)** and how many species are represented by **2 individuals (doubletons)**:

$$S_{\text{est}} = S_{\text{obs}} + F^2/2G$$

S_{est} (nb of species we want to estimate), S_{obs} (nb of species observed), F (nb of singletons) and G (nb of doubletons)

If the **chao1 is close to the richness** → the part of the missed ASVs is low → the sequencing depth is good.

α -diversity: Chao1

Example of a abundance table, after FROGS processing, with ASVs filtering with 0.005% threshold:

observation_name	observation_sum	complexe-ADN-1	echantillon1-1	echantillon1-2	echantillon1-3	echantillon2-1	echantillon2-2	echantillon2-3
Cluster_1	298637	56	227	234	120	36754	59089	56534
Cluster_2	155012	688	20604	38077	45508	8417	10464	10655
Cluster_3	52753	2469	14	76	68	37	8	19
Cluster_4	34062	3459	5041	11458	12799	0	37	84
Cluster_5	30263	3	10	13	13	570	806	800
Cluster_6	26805	1301	7	51	35	21	6	16
Cluster_7	25237	1015	7	30	34	16	5	14
Cluster_8	20483	893	6	34	19	18	1	16
Cluster_9	26069	2504	32	60	87	26	7	22
Cluster_10	17383	712	5	23	17	19	8	13
Cluster_11	16674	715	6	27	25	26	2	7
Cluster_12	11420	0	37	76	79	19	24	13
Cluster_13	9414	189	0	24	12	6	0	8
Cluster_14	7972	498	3	7	11	7	3	5
Cluster_15	7267	13	0	19	12	11	2	7
Cluster_16	7131	150	3	8	15	11	0	2
Cluster_17	6407	4953	22	7	1	0	13	4
Cluster_18	6538	28	1	10	18	16	0	6
Cluster_19	5633	3	12	12	45	24	0	3
Cluster_20	5223	183	0	5	12	8	1	1
Cluster_21	4078	12	0	6	9	6	0	4
Cluster_22	4507	0	10	13	20	13	0	2
Cluster_23	4232	3	0	10	8	9	0	4
Cluster_24	3404	160	1	4	6	4	1	0
Cluster_25	3857	1	0	3	6	10	0	2
Cluster_26	2616	1926	16	12	9	2	8	9
Cluster_27	2781	2182	7	2	0	0	6	1


singletons
and
doubletons

→ Chao1 computation possible

α -diversity: Shannon and Inv-Simpson

α -diversity is equivalent to the richness : number of species

Shannon	Inv-Simpson
Evenness of the species abundance distribution	Inverse probability that two sequences sampled at random come from the same species



Interpretation :

15 observed species, but according to Shannon, the uneven community acts like there is 7.85 equally abundant species (5.45 for invSimp)

α -diversity indices

1. Chao1 close to Richness \rightarrow all species have been detected
2. higher Shannon index \rightarrow higher homogeneity \rightarrow greater diversity
3. greater invsimpson index \rightarrow greater diversity

Exploring biodiversity : α -diversity

α -diversity indices available in phyloseq :

- Species **richness** : number of observed ASV
- **Chao1** : number of observed ASV + estimation of the number of unobserved ASV
- **Shannon** entropy / **Jensen** : the width of the ASV relative abundance distribution. Roughly, it reflects our (in)ability to predict ASV of a randomly picked bacteria.
- **Simpson** : $1 -$ probability that two bacteria picked at random in the community belong to different ASV
- **Inverse Simpson** : inverse of the probability that two bacteria picked at random belong to the same ASV
- Other estimators of alpha diversity exist (Chao2, ACE, ICE,...), however the indices presented above allow us to understand alpha diversity with sufficient precision

Exploring biodiversity : α -diversity

FROGSSTAT Phyloseq Alpha Diversity with richness plot
(Galaxy Version 4.1.0+galaxy1)

☆ Favorite Versions ▾ Options

Phyloseq object (format: RData)

4: FROGSSTAT Phyloseq Import Data SUBSAMPLED: asv_data.Rdata

Explore the sample **NORMALISED** count

This file is the result of FROGS Phyloseq Import Data tool

Experiment variable

EnvType

Choose a sample variable to organize graphics test on **EnvType**

The experiment variable that you want to analyse. (--varExp)

The alpha diversity indices to compute

Select/Unselect all

- Observed
- Chao1
- Shannon
- InvSimpson
- Simpson
- ACE
- Fisher

Choose which α -diversity indices you want to compute

(--alpha-measures)

Exercise 5

1. What are the output files ?
2. Which interpretation could you make on the boxplot results ?
3. Does EnvType has an impact on α -diversity indices ?

Exercise 5

1. What are the output files ?

→ Tabular file: contains the detailed value of indices in each sample

→ HTML report: graphical and statistical results

Exercise 5

1. What are the output files ?

→ Tabular file: contains the detailed value of indices in each sample

1	2	3	4	5	6
	Observed	Chao1	se.chao1	Shannon	InvSimpson
BHT0.LOT01	89	90.875	2.25640704112416	2.46283438240559	6.4374614755645
BHT0.LOT03	129	134.2	3.98819923457003	3.01399812576966	11.6378947553209
BHT0.LOT04	137	152	8.65612088483201	2.77419314445453	7.04904738429417
BHT0.LOT05	127	132.526315789474	3.97261840192821	2.82922278153272	7.54330476122993
BHT0.LOT06	135	136	1.30982775947977	2.6365904270666	6.30810073317464
BHT0.LOT07	126	141.260869565217	7.7960250320146	2.36922299088995	5.65591172677601
BHT0.LOT08	172	189.652173913043	8.66767047151361	3.32220303923076	11.229239617499
BHT0.LOT10	155	173.9	9.42281349646639	2.96129964607031	7.55645792419119
CDT0.LOT02	73	87.5263157894737	7.85749286229502	0.968874997875041	1.93691052993399
CDT0.LOT04	145	168.25	10.9999446485673	3.1208274916296	11.0298385276267

Exercise 5


1. What are the output files ?

→ HTML report: graphical and statistical results

Exercise 5

1 dot = 1 sample


One graph per asked indice



Exercise 5

more readable thanks to boxplots

Alpha diversity distribution in function of EnvType




Exercise 5

Same legend for all indices




Scales in y axis are different
(≠ values for each alpha index)

x axis: 8 boxplots for each indices
(4 indices, 8 EnvTypes)



Exercise 5

2. Which interpretation could you make on the boxplot results ?




Exercise 5

2. Which interpretation could you make on the boxplot results ?

- Same image in same scale for Richness and Chao1
→ most species have been detected
- High variability in the number of ASVs per EnvType
- Many taxa observed in **DesLardons**
(highest observed richness)
- Most foods have low effective diversities
(Shannon & InvSimpson)
→ communities are dominated
by few abundant taxa

Alpha diversity distribution in function of EnvType



Exercise 5

Richness plot

Richness plot with boxplot

Alpha Diversity Indices Anova Analysis

Rarefaction curves



3. Does EnvType has an impact on α -diversity indices ?

- What is an ANOVA used for?

→ Test the significance of the previous observations by performing an ANOVA of alpha-diversity

indices against the covariate of interest (EnvType)

Exercise 5

3. Does EnvType has an impact on α -diversity indices ?

Anova interpretations

```
#####
#Perform ANOVA on Observed, which effects are significant
anova.Observed <-aov( Observed ~ Depth + EnvType, anova_data)
summary(anova.Observed)
      Df Sum Sq Mean Sq F value    Pr(>F)
EnvType  7  57656    8237   7.705 1.68e-06 ***
Residuals 56  59864    1069
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#####
#Perform ANOVA on Chaol, which effects are significant
anova.Chaol <-aov( Chaol ~ Depth + EnvType, anova_data)
summary(anova.Chaol)
      Df Sum Sq Mean Sq F value    Pr(>F)
EnvType  7  65691    9384   8.482 4.85e-07 ***
Residuals 56  61954    1106
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

#####
#Perform ANOVA on Shannon, which effects are significant
anova.Shannon <-aov( Shannon ~ Depth + EnvType, anova_data)
summary(anova.Shannon)
      Df Sum Sq Mean Sq F value Pr(>F)
EnvType  7   7.61  1.0866   1.695  0.129
Residuals 56  35.89  0.6409

#####
#Perform ANOVA on InvSimpson, which effects are significant
anova.InvSimpson <-aov( InvSimpson ~ Depth + EnvType, anova_data)
summary(anova.InvSimpson)
      Df Sum Sq Mean Sq F value Pr(>F)
EnvType  7  392.8   56.12   1.261  0.286
Residuals 56 2492.7   44.51
```

Exercise 5

3. Does EnvType has an impact on α -diversity indices ?

Anova interpretations

Does the EnvType have an effect on Observed indice ?

```
#####  
#Perform ANOVA on Observed, which effects are significant  
anova.Observed <-aov( Observed ~ Depth + EnvType, anova_data)  
summary(anova.Observed)  
              Df Sum Sq Mean Sq F value    Pr(>F)      
EnvType      7  57656    8237   7.705 1.68e-06 ***  
Residuals   56  59864    1069  
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#####  
#Perform ANOVA on Chaol, which effects are significant  
anova.Chaol <-aov( Chaol ~ Depth + EnvType, anova_data)  
summary(anova.Chaol)  
              Df Sum Sq Mean Sq F value    Pr(>F)      
EnvType      7  65691    9384   8.482 4.85e-07 ***  
Residuals   56  61954    1106  
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#####  
#Perform ANOVA on Shannon, which effects are significant  
anova.Shannon <-aov( Shannon ~ Depth + EnvType, anova_data)  
summary(anova.Shannon)  
              Df Sum Sq Mean Sq F value    Pr(>F)      
EnvType      7   7.61   1.0866   1.695  0.129  
Residuals   56  35.89   0.6409
```

```
#####  
#Perform ANOVA on InvSimpson, which effects are significant  
anova.InvSimpson <-aov( InvSimpson ~ Depth + EnvType, anova_data)  
summary(anova.InvSimpson)  
              Df Sum Sq Mean Sq F value    Pr(>F)      
EnvType      7  392.8    56.12   1.261  0.286  
Residuals   56 2492.7    44.51
```

Exercise 5

3. Does EnvType has an impact on α -diversity indices ?

Anova interpretations

- Environments differ in terms of richness but not in terms of Shannon and InvSimpson diversity
- This means that all EnvTypes have similar structures (equivalent distributions between several minor ASVs and few dominant ASVs). Even if 2 samples of "Crevette" displayed very high invSimpson (their bacteria were thus more homogeneously distributed), these two samples were not sufficient to make "Crevette" significantly different from the others EnvType.

→ There is no significant difference between the EnvType

```
#####  
#Perform ANOVA on Observed, which effects are significant  
anova.Observed <-aov( Observed ~ Depth + EnvType, anova_data)  
summary(anova.Observed)  
              Df Sum Sq Mean Sq F value    Pr(>F)        
EnvType       7  57656    8237   7.705 1.68e-06 ***  
Residuals    56  59864    1069            
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#####  
#Perform ANOVA on Chaol, which effects are significant  
anova.Chaol <-aov( Chaol ~ Depth + EnvType, anova_data)  
summary(anova.Chaol)  
              Df Sum Sq Mean Sq F value    Pr(>F)        
EnvType       7  65691    9384   8.482 4.85e-07 ***  
Residuals    56  61954    1106            
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#####  
#Perform ANOVA on Shannon, which effects are significant  
anova.Shannon <-aov( Shannon ~ Depth + EnvType, anova_data)  
summary(anova.Shannon)  
              Df Sum Sq Mean Sq F value    Pr(>F)        
EnvType       7   7.61   1.0866   1.695  0.129        
Residuals    56  35.89   0.6409            
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#####  
#Perform ANOVA on InvSimpson, which effects are significant  
anova.InvSimpson <-aov( InvSimpson ~ Depth + EnvType, anova_data)  
summary(anova.InvSimpson)  
              Df Sum Sq Mean Sq F value    Pr(>F)        
EnvType       7   392.8   56.12   1.261  0.286        
Residuals    56 2492.7   44.51            
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Exercise 5

3. Does EnvType has an impact on α -diversity indices ?

Anova interpretations

- Depth does not appear in the results, so there is no effect of depth.
- This is expected as the sequencing depth is equivalent between samples
- If Depth appears as a significant effect, you should normalize

```
#####  
#Perform ANOVA on Observed, which effects are significant  
anova.Observed <-aov( Observed ~ Depth + EnvType, anova_data)  
summary(anova.Observed)  
              Df Sum Sq Mean Sq F value    Pr(>F)      
EnvType       7  57656    8237   7.705 1.68e-06 ***  
Residuals    56  59864    1069  
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```


```
#####  
#Perform ANOVA on Chaol, which effects are significant  
anova.Chaol <-aov( Chaol ~ Depth + EnvType, anova_data)  
summary(anova.Chaol)  
              Df Sum Sq Mean Sq F value    Pr(>F)      
EnvType       7  65691    9384   8.482 4.85e-07 ***  
Residuals    56  61954    1106  
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#####  
#Perform ANOVA on Shannon, which effects are significant  
anova.Shannon <-aov( Shannon ~ Depth + EnvType, anova_data)  
summary(anova.Shannon)  
              Df Sum Sq Mean Sq F value    Pr(>F)      
EnvType       7   7.61   1.0866   1.695  0.129  
Residuals    56  35.89   0.6409
```

```
#####  
#Perform ANOVA on InvSimpson, which effects are significant  
anova.InvSimpson <-aov( InvSimpson ~ Depth + EnvType, anova_data)  
summary(anova.InvSimpson)  
              Df Sum Sq Mean Sq F value    Pr(>F)      
EnvType       7  392.8    56.12   1.261  0.286  
Residuals    56 2492.7    44.51
```

Exercise 5


Rarefaction curve interpretations



Exercise 5

Rarefaction curve interpretations

- Most of the curves reach a plateau
- A deeper sequencing doesn't add more ASVs
- DesLardons reach the plateau later which correspond to a higher Observed




IV. Biodiversity analysis

β -DIVERSITY INDICES

Exploring biodiversity : β -diversity

Many diversity indices are available with the Phyloseq package through the generic distance function.

Different dissimilarities capture different features of the communities.



Exploring biodiversity : β -diversity


There are different ways to measure beta diversity on a dataset, which give different results.

In this example, 3 ways :

- qualitatively, communities are very similar
- quantitatively, communities are very different
- phylogenetically, two communities seem to be closer than the third one.

Which distance to choose?

- No wrong answer. Each beta-diversity indices will characterize communities differently



Exploring biodiversity : β -diversity

If we compare 2 communities A and B:

Jaccard index:


- Fraction of species specific to either A or B → qualitative index

Bray-Curtis index:

- Fraction of the community specific to either A or B → quantitative index

Exploring biodiversity : β -diversity

- 2 communities, Red and Blue
- 15 ASVs with different abundances in Red community and Blue community





Exploring biodiversity : β -diversity

Jaccard index:

- Proportion of species/ASVs specific to either Red or Blue
→ qualitative index
- Pink = common ASVs between the 2 communities (5)
- Red= ASVs specific to Red community (5)
- Blue= ASVs specific to Blue community (5)

$$D_{jac} = 10/15 = 0.667$$





Exploring biodiversity : β -diversity

Bray-Curtis index:

- Proportion of the abundance specific to either Red or Blue → quantitative index
- Ration (sum of specific abundances)/ (total abundances)
- 1st ASV does not contribute (same abundance for Red and Blue communities)
- ASV 2, 3, 4 and 5 contribute up to the excess in one of the communities (8+8+3+3+10) in the sum of specific abundances (Pink is not taken into account in this sum)

$$D_{bc} = (8+8+3+3+10) / (24+26+28+17+9+10) = 0.281$$




Exploring biodiversity : β -diversity

Indices comparison with different distributions:


- between Red & Blue1 communities

- between Red & Blue2 communities




Exploring biodiversity : β -diversity

Jaccard and Bray-Curtis indices are calculated by pairs (in french “deux-à-deux”) so we here compare pair X indices with pair Y indices



Exploring biodiversity : β -diversity

1. Jaccard indices of X and Y are identical \rightarrow same specific fraction (there are as many ASVs specific to Red or Blue1 in X, as there are ASVs specific to Red or Blue2 in Y).
2. Pair X: Bray-Curtis index is low because shared ASVs between Red and Blue1 communities are abundant and specific ASVs are at low abundance.
3. Pair Y: Bray-Curtis index is high because ASVs specific to Red or Blue2 are abundant and shared ASVs are at low abundance




Exploring biodiversity : β -diversity

3 ways to measure beta diversity with the same data set
→ 3 different results.

In this example :

- ✓ qualitatively, communities are very similar
- ✓ quantitatively, communities are very different
- **phylogenetically**, two communities seem to be closer than the third one.




Exploring biodiversity : β -diversity

Unifrac index:

- Fraction of the tree specific to either A or B

Weighted-Unifrac index :

- Fraction of the diversity specific to either A or B



Exploring biodiversity : β -diversity

Unifrac index:

- Fraction of the tree specific to either A or B

$$Unifrac = \frac{\sum \text{specific_branch_length}}{\sum \text{all_branch_length}}$$



3 ASVs identified by sequencing: ASV3, ASV4 in community A and ASV1, ASV3 in community B

Exploring biodiversity : β -diversity

Unifrac index:

- Fraction of the tree specific to either A or B

$$Unifrac = \frac{\sum \text{specific_branch_length}}{\sum \text{all_branch_length}}$$



ASV1 and ASV4 are specific, ASV3 is shared in the 2 communities and ASV2 are absent in the 2 communities

Exploring biodiversity : β -diversity


Unifrac index:

- Fraction of the tree specific to either A or B

If all branch lengths are equal to 1, only branches present in at least one community are taken into account :

$$Unifrac = \frac{\sum \text{specific_branch_length}}{\sum \text{all_branch_length}} = 3/5 = 0.6$$

- Pink = common ASVs between the 2 communities
- Red = tree branch specific to A
- Blue = tree branch specific to B





A reduced branch is a branch whose distance is weighted by the relative abundance of the ASV

Weighted-Unifrac index:

- Fraction of the diversity specific to either A or B

$$WUnifrac = \frac{\sum \text{reduced_branch_length}}{\sum \text{non_reduced_branch_length}}$$





Exploring biodiversity : β -diversity

Weighted-Unifrac index:

- Fraction of the diversity specific to either A or B

$$WUnifrac = \frac{\sum \text{reduced_branch_length}}{\sum \text{non_reduced_branch_length}}$$




Here the specific ASVs (ASV1 and ASV4) are the most abundant and are also the most phylogenetically distant.

Exploring biodiversity : β -diversity


Weighted-Unifrac index:

- Fraction of the diversity specific to either A or B



ratio of the abundance of each branch

$$WUnifrac = \frac{\sum \text{reduced_branch_length}}{\sum \text{non_reduced_branch_length}}$$




A reduced branch is a branch whose distance is weighted by the relative abundance of the ASV

Exploring biodiversity : β -diversity

Weighted-Unifrac index:

- Fraction of the diversity specific to either A or B



$$WUnifrac = \frac{\sum \text{reduced_branch_length}}{\sum \text{non_reduced_branch_length}}$$

$$\text{Blue branches} = \frac{|0 - 0,7|}{|0 + 0,7|} + \frac{|0 - 0,7|}{|0 + 0,7|} = 1 + 1 = 2$$

$$\text{Red branches} = \frac{|0 - 0,8|}{|0 + 0,8|} = 1$$


$$\text{Pink branches} = \frac{|1 - 0,3|}{|1 + 0,3|} + \frac{|0,2 - 0,3|}{|0,2 + 0,3|} = \frac{0,7}{0,3} + \frac{0,1}{0,5} = 0,73$$

$$\sum \text{reduced branch length} = 3,73$$

Exploring biodiversity : β -diversity

Weighted-Unifrac index:

- Fraction of the diversity specific to either A or B



$$WUnifrac = \frac{\sum \text{reduced_branch_length}}{\sum \text{non_reduced_branch_length}}$$

$$\sum \text{non reduced branch length} = 5$$

$$WUnifrac = \frac{\sum \text{reduced_branch_length}}{\sum \text{non_reduced_branch_length}} = \frac{3,73}{5} = 0,75$$

Exploring biodiversity : β -diversity in brief

qualitative indices: presence/absence regardless of abundance

quantitative indices: compare differences in abundance of ASVs

phylogenetic indices: integrate phylogenetic information to qualitative or quantitative indices (weighted or unweighted indices)

Bray-Curtis index : to evaluate the dissimilarity between two given samples, in terms of abundance of ASVs present in each sample. When Bray-Curtis index close to 0 means abundant ASVs are shared and in the same quantities between communities.

Jaccard index: beta diversity index, qualitative, takes into account the fraction of specific ASVs

Unifrac index: beta diversity index, qualitative, takes into account the fraction of specific phylogenetic branches

Weighted-Unifrac index: beta diversity index, quantitative, takes into account the relative abundance of ASVs shared between samples




Exploring biodiversity : β -diversity

→ What do you conclude in terms of Jaccard, Bray Curtis, Unifrac and weighed Unifrac values for these 4 pairs of communities?

 : in common


 : specific to A

 : specific to B



Exploring biodiversity : β -diversity


→ What do you conclude in terms of Jaccard, Bray Curtis, Unifrac and weighed Unifrac values?



High Jaccard: same amount of specific ASVs
Low Unifrac: small distance between specific branches

Exploring biodiversity : β -diversity

→ What do you conclude in terms of Jaccard, Bray Curtis, Unifrac and weighed Unifrac values?




High Jaccard: all ASVs are specific to A or B

High Unifrac: all the branches are specific to A or B

Exploring biodiversity : β -diversity

→ What do you conclude in terms of Jaccard, Bray Curtis, Unifrac and weighed Unifrac values?




High Bray-Curtis: ASVs are shared but abundant ASVs are not the same in each community

Low weighted-Unifrac: abundant ASVs in a community have a phylogenetically close relative in the other community

Exploring biodiversity : β -diversity

→ What do you conclude in terms of Jaccard, Bray Curtis, Unifrac and weighed Unifrac values?



High Bray-Curtis: ASVs are shared but abundant ASVs are not the same in each community

High weighted-Unifrac: abundant ASVs in a community are phylogenetically distant to any ASV in the other community

Exploring biodiversity : β -diversity

Phyloseq supports currently 43 beta diversity distance methods,
(see [phyloseq distanceMethodList documentation](#))

unifrac, wunifrac,

dpcoa, jsd, manhattan, euclidean, canberra,

bray, kulczynski, jaccard, gower, altGower, morisita, horn, mountford, raup, binomial
chao, cao...

Exploring biodiversity : β -diversity

FROGSSTAT Phyloseq Beta Diversity distance matrix (Galaxy
Version 4.1.0+galaxy1)

☆ Favorite

🔄 Versions

▼ Options

Phyloseq object (format: RData)

4: FROGSSTAT Phyloseq Import Data SUBSAMPLED: asv_data.Rdata

This is the result of FROGS Phyloseq Import Data tool.

Grouping variable

EnvType

Experimental variable used to group samples (Treatment, Host type, etc) (--varExp)

The methods of beta diversity

Select/Unselect all

Unifrac
 Weighted Unifrac
 Bray-Curtis
 Jaccard (as cc method in betadiver vegan funcion)

N.B. if the tree is not available in your RData, you cannot choose Unifrac or Weighted Unifrac (--distance-methods)

Other method

The other methods of beta diversity that you want to use (comma separated value). c.f. details below.

Explore the sample **NORMALISED** count

Choose a sample variable to organize graphics.

Choose which beta diversity distances you want to compute

You can ask another beta-diversity method

Exercise 6

Try it with the 4 most commonly used distance methods

1. What are the output datasets ?
2. *A priori*, abundant ASVs are they shared among samples?
3. Comparing Jaccard and Unifrac, what can you conclude ?
4. Comparing Unifrac and weighted Unifrac, what can you conclude ?

Exercise 6

1. What are the output files ?

→ Tabular file: a tabular file per distance method containing the “all samples against all” matrix of beta diversity distance

→ HTML report: heatmap representing the distance matrix computed

FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html (wunifrac.tsv)

FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html (unifrac.tsv)

FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html (cc.tsv)

FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html (bray.tsv)

FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html


For Jaccard




Exercise 6

[FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html](#)


Heatmap plot of the beta distance : bray





Heatmap plot of the beta distance : cc



Heatmap plot of the beta distance : unifrac




Heatmap plot of the beta distance : wunifrac




Exercise 6

- Each square represents a comparison between 2 samples
- Lighter means more similar
- The diagonal represents the comparison of a sample with itself
- Along the diagonal we can spot clearer square structures
- We can assume that these are the different EnvTypes as the samples are ordered.


Bray-Curtis



Jaccard




Heatmap plot of the beta distance : unifrac



Unifrac

Heatmap plot of the beta distance : wunifrac




Weighted-Unifrac

Exercise 6

2. *A priori*, are abundant ASV shared among samples ?


Bray-Curtis

Heatmap plot of the beta distance : Bray




Jaccard

Heatmap plot of the beta distance : cc




Heatmap plot of the beta distance : unifrac



Unifrac

Heatmap plot of the beta distance : wunifrac



Weighted-Unifrac

Exercise 6

2. *A priori*, are abundant ASV shared among samples ?


- Jaccard lower than Bray-Curtis
- Weighted-Unifrac is lower than Unifrac

→ The abundance accentuates the differences i.e. the distances are greater, i.e. the images are darker

→ abundant ASVs are community specific


Bray-Curtis

Heatmap plot of the beta distance : bray




Jaccard

Heatmap plot of the beta distance : cc




Heatmap plot of the beta distance : unifrac



Unifrac

Heatmap plot of the beta distance : wunifrac




Weighted-Unifrac

Exercise 6

3. Comparing Jaccard and Unifrac, what can you conclude ?


Bray-Curtis

Heatmap plot of the beta distance : bray




Jaccard

Heatmap plot of the beta distance : cc




Heatmap plot of the beta distance : unifrac



Unifrac

Heatmap plot of the beta distance : wunifrac




Weighted-Unifrac

Exercise 6

3. Comparing Jaccard and Unifrac, what can you conclude ?

- Jaccard and Unifrac are close.
- the phylogenetic distances do not accentuate the qualitative data of the Jaccard (neither darker, nor lighter), the species are thus close
- ASVs are distinct but phylogenetically related




Exercise 6

4. Comparing Unifrac and weighted Unifrac, what can you conclude ?


Bray-Curtis

Heatmap plot of the beta distance : bray




Jaccard

Heatmap plot of the beta distance : cc




Heatmap plot of the beta distance : unifrac



Unifrac

Heatmap plot of the beta distance : wunifrac



Weighted-Unifrac


Exercise 6

4. Comparing Unifrac and weighted Unifrac, what can you conclude ?

- Unifrac higher/darker than weighted Unifrac so distance between samples are more important
 - taking into account the abundances makes the samples less distant (lighter)
- abundant ASVs in both communities are phylogenetically closed.


Bray-Curtis

Heatmap plot of the beta distance : bray




Jaccard

Heatmap plot of the beta distance : cc




Heatmap plot of the beta distance : unifrac



Unifrac

Heatmap plot of the beta distance : wunifrac



Weighted-Unifrac

Exploring biodiversity : β -diversity

- In general, **qualitative** diversities (Jaccard, Unifrac) **are more sensitive to factors that affect presence/absence** of organisms (such as pH, salinity, depth, etc) and therefore are useful to study and define bioregions (regions with little or no flow between them)...
- ... whereas **quantitative** distances (Bray-Curtis, weighted-Unifrac) **focus on factors that affect relative changes** (seasonal changes, nutrient availability, concentration of oxygen, depth, etc.) and therefore useful to monitor communities over time or along an environmental gradient.

Different distances capture different features of the samples.

There is no "one size fits all"

Exploring the structure

We will try to identify structures, relationships between samples related to environmental factors

I. Structure Visualisation

ORDINATION AND HEATMAP PLOTS

We have calculated distances now, we will use ordination methods to explore them.

Structure visualization : with PCA ?

- Each community can be described by its ASV abundances, which could be used for a PCA, but high number of ASV make interpretations difficult
- Moreover, PCA maximizes variance and can therefore emphasize differences of rare ASVs between samples instead of giving a good representation of resemblances.
Variance is not a very good measure of β -diversity.
- PCA is not design to use diversity indices and/or distances as it requires independency between variables and does not fit to distance matrix, which is not constructed with samples and variables.
 β -diversity indices thus required dedicated PCA-like methods.

Purpose of the tool : ordinate samples based on β -diversity indices and offer tools to visualize it: produce *ordination plots* and *heatmaps*.

Structure visualization : Ordination plot


The Multidimensional Scaling (**MDS** or **PCoA**) is equivalent to a Principal Component Analysis (PCA) but preserves the β -diversity instead of the variance.

The MDS tries to represent samples in two dimensions while preserving the distances

1- calculation of a distance matrix.

Distance Matrix					
	S1	S2	S3	S4	S5
S1	0.00	2.21	6.31	0.99	7.50
S2	2.21	0.00	5.40	1.22	5.74
S3	6.31	5.40	0.00	5.75	3.16
S4	0.99	1.22	5.75	0.00	6.64
S5	7.50	5.74	3.16	6.64	0.00

2- The samples ordination :



Structure visualization : Heatmap

- Heatmap is an other representation of the abundance table.
- It tries to reveal if there is a structure between a group of ASVs and a group of samples.
- Heatmap
 - Finds a meaningful order of the samples and the ASVs
 - Allows the user to choose a custom order (in R)
 - Allows the user to change the colour scale (in R)
 - Produces a ggplot2 object, easy to manipulate and customize

Structure visualization : Ordination plot and Heatmap

FROGSSTAT Phyloseq Structure Visualisation with heatmap plot and ordination plot (Galaxy Version 4.1.0+galaxy1)

☆ Favorite

🔄 Versions

▼ Options

Phyloseq object (format rdata)

4: FROGSSTAT Phyloseq Import Data SUBSAMPLED: asv_data.Rdata

This is the result of FROGS Phyloseq Import Data Tool.

The beta diversity distance matrix file

11: FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html (cc.tsv)

This file is the result of FROGS Phyloseq Beta Diversity tool (--distance-matrix)

Experiment variable

EnvType

The experiment variable that you want to analyse. (--varExp)

Ordination method

MDS/PCoA

(--ordination-method)

Explore the sample **NORMALISED** count

To see all, launch **once per distance to ordinate** (Bray, Jaccard, Unifrac and Weighted-Unifrac matrices)

Choose a sample variable to organize graphics

Choose the ordination method (most commonly used is MDS/PCoA)

Structure visualization : Ordination plot and Heatmap


Try it with the 4 distance matrices

1. What are the output datasets ?
2. What is the best distance matrix to use to better separate samples ?
3. Guess why Lardon are somewhere between Meat and Seafood ?
4. Based on your favourite distance matrix, what can you conclude on the heatmap ?

Structure visualization : Ordination plot and Heatmap

1. What are the output datasets ?


→ HTML report: ordination plot




Structure visualization : Ordination plot and Heatmap

2. What is the best distance matrix to use to better separate samples ?

JACCARD




UNIFRAC




EnvType

- BoeufHache
- Crevette
- DesLardons
- FiletCabillaud
- FiletSaumon
- MerguezVolaille
- SaumonFume
- VeauHache




BRAY


WUNIFRAC




JACCARD




UNIFRAC




BRAY




WUNIFRAC



JACCARD



UNIFRAC





- Qualitative distances (Unifrac, Jaccard) separate meat products from seafood ones
- ➔ detected taxa segregate by origin

Structure visualization : Ordination plot and Heatmap

3. Guess why Lardon are somewhere between Meat and Seafood ?


JACCARD




UNIFRAC

Structure visualization : Ordination plot and Heatmap

3. Guess why Lardon are somewhere between Meat and Seafood ?



JACCARD



UNIFRAC

■ DesLardons is somewhere in between

➔ contamination induced by sea salt

Structure visualization : Ordination plot and Heatmap


Other conclusions ?

1. Quantitative distances (weighted Unifrac) exhibit a 'meat – seafood' gradient (on axis 1) with DesLardons in the middle and a 'SaumonFume - everything else' gradient on axis 2.


Structure visualization : Ordination plot and Heatmap

Other conclusions ?

- Note the difference between weighted-UniFrac and Bray-Curtis (2 quantitative indices) for the distances between BoeufHache and VeauHache.



Very different visualization




BRAY

WUNIFRAC

Structure visualization : Ordination plot and Heatmap

Other conclusions ?

3. On Bray-Curtis, on axis 2, we can observe the distribution of Saumon Fumé samples. Axis 1 shows the distribution of MerguezdeVolaille samples



BRAY

Structure visualization : Ordination plot and Heatmap

Other conclusions ?



The 2D representation captures only parts of the original distances

Ellipse are not always an advantage for visualization because it accentuates the 2D effect

Structure visualization : Ordination plot and Heatmap

4. Based on your favourite distance matrix, what can you conclude on the **heatmap** ?


Try to identify:

- Block-like structure of the abundance table
- Interaction between (groups of) taxa and (groups of) samples
- Core and condition-specific microbiota

Exercise 7


4. Based on your favourite distance matrix, what can you conclude on the heatmap ?

matrix based on Jaccard distance (qualitative) which "sorts" the ASVs. Then a color is applied according to the abundance of ASVs (yellow to red).




Exercise 7

4. Based on your favourite distance matrix, what can you conclude on the heatmap ?



Exercise 7

4. Based on your favourite distance matrix, what can you conclude on the heatmap ?



Note: no evidence for seafood.

II. Exploring the structure


HIERARCHICAL CLUSTERING

Exploring the structure : clustering

Clustering aims to represent samples in a tree based on a distance matrix and a linkage function:

3 clustering algorithms:

- **Complete linkage:** tends to produce compact, spherical clusters and guarantees that all samples in a cluster are similar to each other.
- **Ward:** tends to also produce spherical clusters but has better theoretical properties than complete linkage.
- **Single:** friend of friend approach, tends to produce banana-shaped or chains-like clusters.



Exploring the structure : clustering

FROGSSTAT Phyloseq Sample Clustering of samples using different linkage methods (Galaxy Version 4.1.0+galaxy1)

☆ Favorite

🔄 Versions

▼ Options

Phyloseq object (format: RData)

4: FROGSSTAT Phyloseq Import Data SUBSAMPLED: asv_data.Rdata

Explore the sample **NORMALISED** count

This is the result of FROGS Phyloseq Import Data tool.

The beta diversity distance matrix file

11: FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html (cc.tsv)

Choose the beta diversity distance matrix: i.e. Jaccard

This file is the result of FROGS Phyloseq Beta Diversity tool. (--distance-matrix)

Experiment variable

EnvType

Choose a sample variable to organize graphics: i.e. EnvType

The experiment variable that you want to analyse. (--varExp)

The three different linkage functions will be used, generating three different dendrograms


Exercise 8

Try it with « a good » distance method matrix on EnvType and on FoodType


→ Which linkage method seems to better fit the data ?

Exercise 8


single linkage clustering tree



ward.D2 linkage clustering tree



complete linkage clustering tree




the Ward clustering allows to classify the communities according to the EnvType groups

Exercise 8

- Consistently, for these datasets, with the ordination plots, clustering works quite well for the **UniFrac** distance
- The method (Ward.D2) give almost a perfect separation between the different type of food

Remarks

Clustering is based on the whole distance whereas ordination represents parts of the distance (the most it can with 2 dimensions)



Exercise 8

Ward D2

Complete

Single


ward.D2 linkage clustering tree

- Not as well clustered with **Jaccard** indices
- DesLardons is in the middle of seafood.

Once again,

Different distances capture different features
of the samples.

There is no "one solution fits all"



■ DesLardons	■ BoeufHache	■ SaumonFume	■ FiletCabillaud
■ MerguezVolaille	■ VeauHache	■ FiletSaumon	■ Crevette


Diversity partitioning

Diversity partitioning

Do the structures seem linked to metadata ? Does the metadata have an effect on the composition of our communities ?

To answer these questions, **multivariate analyses** :

- test **composition differences** of communities from different groups **using a distance matrix**
- compare **within-group** to **between-group** distances




Diversity partitioning : Multivariate ANOVA

Idea : Test **differences** in the community composition **from different groups** using a **distance matrix**.

How it works ?

- Computes sum of square distance
- Variance analysis



Diversity partitioning : Multivariate ANOVA

FROGSSTAT Phyloseq Multivariate Analysis Of Variance perform
Multivariate Analysis of Variance (MANOVA) (Galaxy Version 4.1.0+galaxy1)

☆ Favorite

▼ Options

Phyloseq object (format: RData)

69: FROGSSTAT Phyloseq Import Data: asv_data.Rdata

This is the result of FROGS Phyloseq Import Data tool.

The beta diversity distance matrix file

76: FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html (cc.tsv)

This file is the result of FROGS Phyloseq Beta Diversity tool (--distance-matrix)

Experiment variable

EnvType

The experiment variable that you want to analyse (--varExp)

Explore the sample **NORMALISED** count

Choose the beta diversity distance matrix: Unifrac

Choose the variable to explain the variability between samples: EnvType

To simultaneously test several variables, you can use “+” symbol as “EnvType+FoodType” to test only additive effects or “*” symbol as “EnvType*FoodType” to test for additive effects and interactions between variables

Exercise 9

Try it with a good beta distance matrix with EnvType and FoodType

1. Does EnvType have an influence on the beta diversity variance ?
2. What about FoodType ?

Exercise 9

1. Does EnvType have an influence on the beta diversity variance ?

With Unifrac distance

```
Call:
adonis(formula = dist ~ EnvType, data = metadata, permutations = 9999)

Permutation: free
Number of permutations: 9999

Terms added sequentially (first to last)

              Df SumsOfSqs MeanSqs F.Model      R2 Pr(>F)
EnvType       7     6.1849 0.88356  11.164 0.58255 1e-04 ***
Residuals    56     4.4320 0.07914           0.41745
Total        63    10.6170           1.00000
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Exercise 9

1. Does EnvType have an influence on the beta diversity variance ?

Environment type explains roughly **58%** of the total variation, which is very high

With Unifrac distance

```
Call:
adonis(formula = dist ~ EnvType, data = metadata, permutations = 9999)

Permutation: free
Number of permutations: 9999

Terms added sequentially (first to last)

      Df SumsOfSqs MeanSqs F.Model    R2 Pr(>F)
EnvType  7    6.1849  0.88356  11.164 0.58255 1e-04 ***
Residuals 56    4.4320  0.07914    0.41745
Total    63   10.6170          1.00000
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Exercise 9

2. What about FoodType ?

```
Call:
adonis(formula = dist ~ FoodType, data = metadata, permutations = 9999)
```

```
Permutation: free
Number of permutations: 9999
```

With Unifrac distance

```
Terms added sequentially (first to last)
```

	Df	SumsOfSqs	MeanSqs	F.Model	R2	Pr(>F)	
FoodType	1	1.7858	1.78579	12.537	0.1682	1e-04	***
Residuals	62	8.8312	0.14244		0.8318		
Total	63	10.6170			1.0000		

```
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Exercise 9

2. What about FoodType ?

Food type explains only **17 %** of the total variation

With Unifrac distance

```
Call:
adonis(formula = dist ~ FoodType, data = metadata, permutations = 9999)
```

```
Permutation: free
Number of permutations: 9999
```

```
Terms added sequentially (first to last)
```

	Df	SumsOfSqs	MeanSqs	F.Model	R2	Pr(>F)
FoodType	1	1.7858	1.78579	12.537	0.1682	1e-04 ***
Residuals	62	8.8312	0.14244		0.8318	
Total	63	10.6170			1.0000	

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Differential abundance analysis

Differential abundance analysis

Are there ASV with differential abundance between 2 conditions ? And which are they ?

To answer these questions, we perform a differential abundance analysis using DESeq2 on the phyloseq object

The package DESeq2 provides methods to test for differential expression by use of negative binomial generalized linear models

Differential abundance analysis

Are there ASV with differential abundance between 2 conditions ? And which are they ?

To answer these questions, we perform a differential abundance analysis using DESeq2 on the phyloseq object

The package DESeq2 provides methods to test for differential expression by use of negative binomial generalized linear models



Be aware to use data *without normalisation*

DESeq has its own normalisation method suited to this kind of data.

It uses the postcount function optimised for metagenomic count table

Differential abundance analysis

→ 1st step: launch *DESeq2 Preprocess* tool to create the **dds object** – the DESeq2 object

FROGSSTAT DESeq2 Preprocess import a Phyloseq object and prepare it for DESeq2 differential abundance analysis (Galaxy Version 4.1.0+galaxy1)

☆ Favorite

🔄 Versions

▼ Options

Type of analysis

- ASV
 FUNCTION

Type of data to perform the differential analysis. ASV: DESeq2 is run on the ASV abundance table. FUNCTION: DESeq2 is run on predicted function abundance table from FROGSFUNC_2_function tool.

Ask for DESeq2 **ASV** data analysis

Phyloseq object

   19: FROGSSTAT Phyloseq Import Data NOT NORMALISED: asv_data.Rdata  

This is the result of FROGSSTAT_Phyloseq_Import_Data without normalisation (DESeq2 is more powerful on unnormalised counts) (format RData)

Explore the sample **RAW** count

Experimental variable

EnvType

The factor that could have an effect on ASV/FUNCTION abundances. Ex: Treatment, etc.

Choose the factor on which the differential abundances will be compared

Do you want to correct a confounding factor?

False

If yes, specify the confounding factor

Specify a confounding factor if necessary (example : testing antibiotic treatment effect with 2 different mice phenotypes, or testing drought effect on soil microbiome with two soil compositions)

Differential abundance analysis

→ What are the output datasets ?

→ Rdata file: `asv_dds.Rdata` object with results of the DESeq analysis

→ 2nd step: launch *DESeq2 visualization* tool to explore the `dds` object

Differential abundance visualization

FROGSSTAT DESeq2 Visualisation to extract and visualise differentially abundant ASVs or functions (Galaxy Version 4.1.0+galaxy1)

☆ Favorite

🔗 Versions

▼ Options

Type of analysis

- ASV
 FUNCTION

Type of data to perform the differential analysis. ASV: DESeq2 is run on the ASV abundance table. FUNCTION: DESeq2 is run on predicted function abundance table from FROGSFUNC_2_function tool.

Data object (format: data.RData)

   19: FROGSSTAT Phyloseq Import Data NOT NORMALISED: asv_data.Rdata  

For ASV: asv_data.Rdata from FROGSSTAT_Phyloseq_Import_Data tool - For FUNCTION: function_data.Rdata from FROGSSTAT_DESeq2_Preprocess tool. (--abundanceData)

DESeq2 object (format: dds.RData)

   21: FROGSSTAT DESeq2 Preprocess: asv_dds.Rdata  

This is the result of FROGSSTAT_DESeq2_Preprocess tool asv_dds.Rdata or function_dds.Rdata (--dds)

Ask for DESeq2 **ASV data** analysis

Result of FROGSSTAT DESeq2 preprocess

Factor on which the differential abundances have been tested

Differential abundance visualization

Experimental variable

EnvType

The factor that could have an effect on ASV/FUNCTION abundances. Ex : Treatment (var)

The experimental variable is it quantitative or qualitative?

Qualitative

If qualitative, choose 2 conditions to compare

Condition 1 considered as reference

BoeufHache

One condition of the experimental variable (e.g. with) (--mod2)

Condition 2 to be compared to the reference

VeauHache

Another condition of the experimental variable (e.g. without) (--mod1)

Adjusted p-value threshold

0.05

Threshold used for statistical significance of the differentially abundant ASV/FUNCTION analysis (--padj)

Factor on which the differential abundances have been tested

Specify qualitative or quantitative

Precise the two conditions to compare

Statistical significance threshold (default 0.05)

Differential abundance visualization

What are the output datasets ?

→ HTML report: result table and several plots

Differentially abundant ASV/FUNCTION table

Pie chart

Volcano plot

MA plot

Heatmap plot

Differential abundance visualization

Differentially abundant ASV/FUNCTION table

[Pie chart](#)

[Volcano plot](#)

[MA plot](#)

[Heatmap plot](#)

Code

```
You chose to compare VeauHache to the reference modality BoeufHache. This implies that a positive log2FoldChange means more abundant in VeauHache than in BoeufHache.
```

Code

Then we extract significant ASVs or FUNCTIONS at the p-value adjusted threshold (after Benjamini Hochberg correction) and enrich results with taxonomic/functional classification and sort the results by pvalue.

Differential abundance visualization

Download Search:

ID	baseMean	log2FoldChange	lfcSE	stat	pvalue	padj	Kingdom	
<input type="text"/>	A	All	<input type="text"/>	<input type="text"/>	All	All	<input type="text"/>	
1	Cluster_53	16.7845	-7.93954	1.21935	-6.51127	7.45192e-11	2.61563e-8	Bacteria
2	Cluster_43	10.4196	15.6431	2.48659	6.29099	3.15446e-10	5.53607e-8	Bacteria
3	Cluster_120	7.49645	5.21487	0.842194	6.19200	5.94040e-10	6.95027e-8	Bacteria
4	Cluster_4	284.010	-4.46973	0.730032	-6.12265	9.20307e-10	8.07569e-8	Bacteria
5	Cluster_85	5.25312	-14.8545	2.69005	-5.52204	3.35093e-8	0.00000235236	Bacteria
6	Cluster_174	2.99262	-17.3671	3.27384	-5.30481	1.12788e-7	0.00000659810	Bacteria
7	Cluster_44	22.0406	-6.03398	1.14995	-5.24715	1.54472e-7	0.00000677746	Bacteria
8	Cluster_141	9.26135	5.96649	1.13629	5.25083	1.51415e-7	0.00000677746	Bacteria
9	Cluster_9	150.302	28.4432	5.83716	4.87279	0.00000110034	0.0000429134	Bacteria
10	Cluster_135	7.45843	-4.76315	1.05240	-4.52600	0.00000601095	0.000210984	Bacteria

Show entries

Showing 1 to 10 of 35 entries

Previous 2 3 4 Next

Differentially abundant ASV/FUNCTION table

Only significantly differentially abundant ASV are displayed (with an adjusted p-value < previously defined threshold - set here to 0.05)

p-value are adjusted using the Benjamini-Hochberg method

Differential abundance visualization

ID	baseMean	log2FoldChange	lfcSE	stat	pvalue	padj	Kingdom
<input type="text"/>	A	All	<input type="text"/>	<input type="text"/>	All	All	<input type="text"/>
1	Cluster_53	16.7845	-7.93954	1.21935	-6.51127	7.45192e-11	
2	Cluster_43	10.4196	15.6431	2.48659	6.29099	3.15446e-10	5.53607e-8 Bacteria
3	Cluster_120	7.49645	5.21487	0.842194	6.19200	5.94040e-10	6.95027e-8 Bacteria
4	Cluster_4	284.010	-4.46973	0.730032	-6.12265	9.20307e-10	
5	Cluster_85	5.25312	-14.8545	2.69005	-5.52204	3.35093e-8	0.00000235236 Bacteria
6	Cluster_174	2.99262	-17.3671	3.27384	-5.30481	1.12788e-7	0.00000659810 Bacteria
7	Cluster_44	22.0406	-6.03398	1.14995	-5.24715	1.54472e-7	0.00000677746 Bacteria
8	Cluster_141	9.26135	5.96649	1.13629	5.25083	1.51415e-7	0.00000677746 Bacteria
9	Cluster_9	150.302	28.4432	5.83716	4.87279	0.00000110034	0.0000429134 Bacteria
10	Cluster_135	7.45843	-4.76315	1.05240	-4.52600	0.00000601095	0.000210984 Bacteria

Differentially abundant ASV/FUNCTION table

More abundant in BoeufHache than VeauHache

More abundant in VeauHache than BoeufHache

Differential abundance visualization

Why log2Foldchange ?

Differentially abundant ASV/FUNCTION table

Foldchange:

It's the ratio of the normalized counts between VeauHache and BoeufHache

log2 is used for interpret and scale reasons:

- Positive values denote an increase, and negative a decrease of abundance
- $\log_2FC = 1$ means a doubling
- $\log_2FC = 2$ means a quadrupling
- $\log_2FC = -1$ means a halving
- $\log_2FC = -2$ means a quartering
- ...

Differential abundance visualization

ID	baseMean	log2FoldChange	lfcSE	stat	pvalue	padj	Kingdom	
<input type="text"/>	A	All	<input type="text"/>	<input type="text"/>	All	All	<input type="text"/>	
1	Cluster_53	16.7845	-7.93954	1.21935	-6.51127	7.45192e-11	2.61563e-8	Bacteria
2	Cluster_43	10.4196	15.6431	2.48659	6.29099	3.15446e-10	5.53607e-8	Bacteria
3	Cluster_120	7.49645	5.21487	0.842194	6.19200	5.94040e-10	6.95027e-8	Bacteria
4	Cluster_4	284.010	-4.46973	0.730032	-6.12265	9.20307e-10	8.07569e-8	Bacteria
5	Cluster_85	5.25312	-14.8545	2.69005	-5.52204	3.35093e-8	0.00000235236	Bacteria
6	Cluster_174	2.99262	-17.3671	3.27384	-5.30481	1.12788e-7	0.00000659810	Bacteria
7	Cluster_44	22.0406	-6.03398	1.14995	-5.24715	1.54472e-7	0.00000677746	Bacteria
8	Cluster_141	9.26135	5.96649	1.13629	5.25083	1.51415e-7	0.00000677746	Bacteria
9	Cluster_9	150.302	28.4432	5.83716	4.87279	0.00000110034	0.0000429134	Bacteria
10	Cluster_135	7.45843	-4.76315	1.05240	-4.52600	0.00000601095	0.000210984	Bacteria

Differentially abundant ASV/FUNCTION table

You can sort by numeric columns and filter on taxonomy

Differential abundance visualization

→ Which species have the highest positive log2Foldchange ?

Differentially abundant ASV/FUNCTION table

	ID	baseMean	log2FoldChange	lfcSE	stat	pvalue	padj	Kingdom
	<input type="text"/>	<input type="text" value="A"/>	<input type="text" value="All"/>	<input type="text"/>	<input type="text"/>	<input type="text" value="All"/>	<input type="text" value="All"/>	<input type="text"/>
1	Cluster_53	16.7845	-7.93954	1.21935	-6.51127	7.45192e-11	2.61563e-8	Bacteria
2	Cluster_43	10.4196	15.6431	2.48659	6.29099	3.15446e-10	5.53607e-8	Bacteria
3	Cluster_120	7.49645	5.21487	0.842194	6.19200	5.94040e-10	6.95027e-8	Bacteria
4	Cluster_4	284.010	-4.46973	0.730032	-6.12265	9.20307e-10	8.07569e-8	Bacteria
5	Cluster_85	5.25312	-14.8545	2.69005	-5.52204	3.35093e-8	0.00000235236	Bacteria
6	Cluster_174	2.99262	-17.3671	3.27384	-5.30481	1.12788e-7	0.00000659810	Bacteria
7	Cluster_44	22.0406	-6.03398	1.14995	-5.24715	1.54472e-7	0.00000677746	Bacteria
8	Cluster_141	9.26135	5.96649	1.13629	5.25083	1.51415e-7	0.00000677746	Bacteria

Differential abundance visualization

→ Which species have the highest positive log2Foldchange (more present in VeauHache than BoeufHache) ?

ID	baseMean	log2FoldChange	
	A	All	
9	Cluster_9	150.302	28.4432



It's the Cluster_9 which is a *Weissella ceti*

Phylum	Class	Order	Family	Genus	Species
All	All	All	All	All	All
Firmicutes	Bacilli	Lactobacillales	Lactobacillaceae	Weissella	Weissella ceti

Differential abundance visualization

Pie chart to view ASVs or FUNCTIONS number of Differential Abundance test

Pie chart




Most of the ASVs are not significantly affected between the conditions (DESeq2 hypothesis !!)

35 ASVs are significantly affected between conditions

Differential abundance visualization

Volcano Plot
Colored by effect sign

Volcano plot




visualization of ASVs $\log_2\text{FoldChange}$ and their associated adjusted p-values

Only ASVs with a significant adjusted p-value are colored

Differential abundance visualization

Post Normalisation DESeq2: MA plot of log2FoldChange

MA plot



visualization of the relation between log2foldchange between conditions, and mean abundance of ASVs (significantly affected ASVs are colored)

Colored ASVs on the right : abundant ASVs affected by the conditions


Colored ASVs on the left : affected rare ASVs

Triangles represent ASV out of scale

Differential abundance visualization

Post Normalisation DESeq2: MA plot of log2FoldChange


MA plot



→ Which Cluster is the triangle spotted?

Differential abundance visualization

Post Normalisation DESeq2: MA plot of log2FoldChange



MA plot

➔ Which Cluster is the triangle spotted?

It's Cluster_9 !

mean abundance

ID	baseMean	log2FoldChange
	A	All
9	Cluster_9	150.302
		28.4432

Differential abundance visualization


Heatmap plot of DA asv or functions, between 2 conditions
EnvType_VeauHache_vs_BoeufHache

Heatmap plot



visualization of the DESeq2 normalised abundances of differentially abundant ASVs grouped by condition

Here, we observe only the significant 35 ASV that are differential abundant



ASVs are ordered from top to bottom in log2 fold change descending order

Differential abundance visualization

Compare FiletSaumon vs SaumonFume

Experimental variable

EnvType

The factor that could have an effect on ASV/FUNCTION abundances. Ex : Treatment (var)

The experimental variable is it quantitative or qualitative?

Qualitative

If qualitative, choose 2 conditions to compare

Condition 1 considered as reference

FiletSaumon

One condition of the experimental variable (e.g. with) (--mod2)

Condition 2 to be compared to the reference

SaumonFume

Another condition of the experimental variable (e.g. without) (--mod1)

Differential abundance visualization

Differentially abundant ASV/FUNCTION table

[Pie chart](#)

[Volcano plot](#)

[MA plot](#)

[Heatmap plot](#)

Since we only have a binary factor we can use the following syntax to format the log₂ fold change from the fitted model if not, we will use the other syntax with contrast=c()

Code

```
You chose to compare SaumonFume to the reference modality FiletSaumon. This implies that a positive log2FoldChange means more abundant in SaumonFume than in FiletSaumon.
```

Then we extract significant OTUs at the p-value adjusted threshold level (after correction) and enrich results with taxonomic informations and sort taxa by pvalue.

Differential abundance visualization


	ID	baseMean	log2FoldChange	lfcSE	stat	pvalue	padj	Kingdom
	<input type="text"/>	<input type="text" value="A"/>	<input type="text" value="All"/>	<input type="text"/>	<input type="text"/>	<input type="text" value="All"/>	<input type="text" value="All"/>	<input type="text"/>
1	Cluster_4	284.010	-4.97034	0.718373	-6.91888	4.55218e-12	2.25333e-9	Bacteria
2	Cluster_85	5.25312	-17.5013	2.66091	-6.57717	4.79475e-11	1.18670e-8	Bacteria
3	Cluster_55	19.0634	-4.83859	0.825830	-5.85906	4.65500e-9	7.68076e-7	Bacteria
4	Cluster_123	10.3886	7.90236	1.39576	5.66171	1.49873e-8	0.00000185468	Bacteria
5	Cluster_31	37.4358	-5.51672	1.04587	-5.27478	1.32918e-7	0.0000131588	Bacteria
6	Cluster_13	139.041	4.03643	0.838190	4.81565	0.00000146724	0.000121047	Bacteria
7	Cluster_27	41.5512	-5.32505	1.13155	-4.70599	0.00000252641	0.000178653	Bacteria

Differentially abundant ASV/FUNCTION table

Differential abundance visualization


Pie chart to view OTUs number of Differential Abundance test

Pie chart



Most of the ASV are not significantly affected between your conditions


Only 47 ASVs are significantly affected between conditions



Differential abundance visualization


Volcano Plot
Colored by effect sign

Volcano plot



→ Which Cluster is it ?

Differential abundance visualization



Volcano plot

→ Which Cluster is it ?


Cluster_85: *Flavobacterium omnivorum*

	ID	baseMean	log2FoldChange
	<input type="text"/>	<input type="text" value="A"/>	<input type="text" value="All"/>
2	Cluster_85	5.25312	-17.5013
22	Cluster_76	12.5611	-8.43272
9	Cluster_73	7.76604	-6.95033

Differential abundance visualization

Post Normalisation DESeq2: MA plot of log2FoldChange


MA plot



Differential abundance visualization

Heatmap plot of DA asv or functions, between 2 conditions

EnvType_SaumonFume_vs_FiletSaumon



Heatmap plot

FROGSStat Summary

FROGSSTAT
Phyloseq Import Data

- Abundance biom file with taxonomical metadata (format: BIOM)
- Metadata associated to samples (format: TSV)
- Taxonomic tree file (format: Newick)
- FROGSSTAT Phyloseq Import Data: asv_data.Rdata (rdata)
- FROGSSTAT Phyloseq Import Data: report.nb.html (html)

FROGSSTAT
Phyloseq Composition Visualisation

- Phyloseq object (format: rdata)
- FROGSSTAT Phyloseq Composition Visualisation: composition.nb.html (html)

FROGSSTAT
Phyloseq Alpha Diversity

- Phyloseq object (format: RData)
- FROGSSTAT Phyloseq Alpha Diversity: alpha_diversity.tsv (tsv)
- FROGSSTAT Phyloseq Alpha Diversity: alpha_diversity.nb.html (html)

FROGSSTAT
Phyloseq Beta Diversity

- Phyloseq object (format: RData)
- FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html (html)

What is the sample composition ?

What are the sample diversities ?

Composition analysis

Structure analysis

Is there any relation between species or communities?

How do the communities cluster?

Which variable influence the diversity ?

What is the samples dissimilarity ?

Differential analysis

Which ASVs are differentially abundant?

FROGSSTAT
DESeq2 Preprocess

- Phyloseq object
- FROGSSTAT DESeq2 Preprocess: asv_dds.Rdata (rdata)

FROGSSTAT
DESeq2 Visualisation

- Data object (format: data.RData)
- DESeq2 object (format: dds.RData)
- FROGSSTAT DESeq2 Visualisation: report.nb.html (html)

FROGSSTAT
Phyloseq Structure Visualisation

- Phyloseq object (format: rdata)
- The beta diversity distance matrix file
- FROGSSTAT Phyloseq Structure Visualisation: structure.nb.html (html)

FROGSSTAT
Phyloseq Sample Clustering

- Phyloseq object (format: RData)
- The beta diversity distance matrix file
- FROGSSTAT Phyloseq Sample Clustering: clustering.nb.html (html)

FROGSSTAT
Phyloseq Multivariate Analysis Of Variance

- Phyloseq object (format: RData)
- The beta diversity distance matrix file
- FROGSSTAT Phyloseq Multivariate Analysis Of Variance: manova.nb.html (html)

FROGSStat Summary

FROGSSTAT Phyloseq Import Data

- Abundance biom file with taxonomical metadata (format: BIOM)
- Metadata associated to samples (format: TSV)
- Taxonomic tree file (format: Newick)
- FROGSSTAT Phyloseq Import Data: asv_data.Rdata (rdata)
- FROGSSTAT Phyloseq Import Data: report.nb.html (html)

FROGSSTAT Phyloseq Composition Visualisation

- Phyloseq object (format: rdata)
- FROGSSTAT Phyloseq Composition Visualisation: composition.nb.html (html)

FROGSSTAT Phyloseq Alpha Diversity

- Phyloseq object (format: RData)
- FROGSSTAT Phyloseq Alpha Diversity: alpha_diversity.tsv (tsv)
- FROGSSTAT Phyloseq Alpha Diversity: alpha_diversity.nb.html (html)

FROGSSTAT Phyloseq Beta Diversity

- Phyloseq object (format: RData)
- FROGSSTAT Phyloseq Beta Diversity: beta_diversity.nb.html (html)

FROGSSTAT Phyloseq Structure Visualisation

- Phyloseq object (format: rdata)
- The beta diversity distance matrix file
- FROGSSTAT Phyloseq Structure Visualisation: structure.nb.html (html)

FROGSSTAT Phyloseq Sample Clustering

- Phyloseq object (format: RData)
- The beta diversity distance matrix file
- FROGSSTAT Phyloseq Sample Clustering: clustering.nb.html (html)

FROGSSTAT Phyloseq Multivariate Analysis Of Variance

- Phyloseq object (format: RData)
- The beta diversity distance matrix file
- FROGSSTAT Phyloseq Multivariate Analysis Of Variance: manova.nb.html (html)

FROGSSTAT DESeq2 Preprocess

- Phyloseq object
- FROGSSTAT DESeq2 Preprocess: asv_dds.Rdata (rdata)

FROGSSTAT DESeq2 Visualisation

- Data object (format: data.RData)
- DESeq2 object (format: dds.RData)
- FROGSSTAT DESeq2 Visualisation: report.nb.html (html)

What is the sample composition ?
What are the sample diversities ?


Composition analysis

Structure analysis

Is there any relation between species or communities?



How do the communities cluster?

Which variable influence the diversity ?



Differential analysis

Which ASVs are differentially abundant?



```

adonis(formula = dist ~ EnvType, data = metadata, permutations = 9999)
Permutation: free
Number of permutations: 9999
Terms added sequentially (first to last)

          Df SumSqs  MeanSS  F.Model    Pr(>F)
EnvType  7  6.1849  0.88356  11.164 0.0255 1e-04 ***
Residuals 56  4.4320  0.07914   0.41745
Total    63 10.6170
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
    
```

Conclusion and advices reminder

FROGSTAT advices

- Before starting, **check taxonomy format** : how many levels? What are their names ?
- Carefully construct your **sample_metadata** TSV file, and after its import, check that your variable order is smart
- Keep in mind that :
 - **Phyloseq composition** and **structure analyses** need to be performed on **normalised** (=rarefied) counts
 - **DESeq** analysis needs to be performed on counts **without normalisation**
 - Different indices or distance methods will give **different but complementary** information
 - **Test different distances and choose which one fits better your data**

References

- Chaillou, S., Chaulot-Talmon, A., Caekebeke, H., Cardinal, M., Christieans, S., Denis, C., Desmonts, M. H., Dousset, X., Feurer, C., Hamon, E., Joraud, J.-J., La Carbona, S., Leroi, F., Leroy, S., Lorre, S., Mace, S., Pilet, M.-F., Prevost, H., Rivollier, M., Roux, D., Talon, R., Zagorec, M., and Champomier-Verges, M.-C. (2015). Origin and ecological selection of core and food-specific bacterial communities associated with meat and seafood spoilage. *ISME J*, 9(5):1105{1118.
- McMurdie, P. J. and Holmes, S. (2013). phyloseq: An r package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE*, 8(4):e61217.
- Shade, A., Jones, S. E., Caporaso, J. G., Handelsman, J., Knight, R., Fierer, N., and Gilbert, J. A. (2014). Conditionally rare taxa disproportionately contribute to temporal changes in microbial diversity. *MBio*, 5(4):e01371{e01314.