



Published in final edited form as:

Nat Methods. 2010 May ; 7(5): 335–336. doi:10.1038/nmeth.f.303.

QIIME allows analysis of high-throughput community sequencing data

J Gregory Caporaso^{1,12}, Justin Kuczynski^{2,12}, Jesse Stombaugh^{1,12}, Kyle Bittinger³, Frederic D Bushman³, Elizabeth K Costello¹, Noah Fierer⁴, Antonio Gonzalez Peña⁵, Julia K Goodrich⁵, Jeffrey I Gordon⁶, Gavin A Huttley⁷, Scott T Kelley⁸, Dan Knights⁵, Jeremy E Koenig⁹, Ruth E Ley⁹, Catherine A Lozupone¹, Daniel McDonald¹, Brian D Muegge⁶, Meg Pirrung¹, Jens Reeder¹, Joel R Sevinsky¹⁰, Peter J Turnbaugh⁶, William A Walters², Jeremy Widmann¹, Tanya Yatsunenکو⁶, Jesse Zaneveld², and Rob Knight^{1,11}

Rob Knight: rob.knight@colorado.edu

¹Department of Chemistry and Biochemistry, University of Colorado, Boulder, Colorado, USA

²Department of Molecular, Cellular and Developmental Biology, University of Colorado, Boulder, Colorado, USA

³Department of Microbiology, University of Pennsylvania, Philadelphia, Pennsylvania, USA

⁴Cooperative Institute for Research in Environmental Sciences and Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, Colorado, USA

⁵Department of Computer Science, University of Colorado, Boulder, Colorado, USA

⁶Center for Genome Sciences, Washington University School of Medicine, St. Louis, Missouri, USA

⁷Computational Genomics Laboratory, John Curtin School of Medical Research, The Australian National University, Canberra, Australian Capital Territory, Australia

⁸Department of Biology, San Diego State University, San Diego, California, USA

⁹Department of Microbiology, Cornell University, Ithaca, New York, USA

¹⁰Luca Technologies, Golden, Colorado, USA

¹¹Howard Hughes Medical Institute, Boulder, Colorado, USA

To the Editor

High-throughput sequencing is revolutionizing microbial ecology studies. Efforts like the Human Microbiome Projects¹ and the US National Ecological Observatory Network² are helping us to understand the role of microbial diversity in habitats within our own bodies and throughout the planet.

Pyrosequencing using error-correcting, sample-specific barcodes allows hundreds of communities to be analyzed simultaneously in multiplex³. Integrating information from

© 2010 Nature America, Inc. All rights reserved.

Correspondence to: Rob Knight, rob.knight@colorado.edu.

¹²These authors contributed equally to this work.

COMPETING FINANCIAL INTERESTS The authors declare competing financial interests: details accompany the full-text HTML version of the paper at <http://www.nature.com/naturemethods/>.

Note: Supplementary information is available on the Nature Methods website.

thousands of samples, including those obtained from time series, can reveal large-scale patterns that were inaccessible with lower-throughput sequencing methods. However, a major barrier to achieving such insights has been the lack of software that can handle these increasingly massive datasets. Although tools exist to perform library demultiplexing and taxonomy assignment^{4,5}, tools for downstream analyses are scarce.

Here we describe ‘quantitative insights into microbial ecology’ (QIIME; pronounced ‘chime’), an open-source software pipeline built using the PyCogent toolkit⁶, to address the problem of taking sequencing data from raw sequences to interpretation and database deposition. QIIME, available at <http://qiime.sourceforge.net/>, supports a wide range of microbial community analyses and visualizations that have been central to several recent high-profile studies, including network analysis, histograms of within- or between-sample diversity and analysis of whether ‘core’ sets of organisms are consistently represented in certain habitats. QIIME also provides graphical displays that allow users to interact with the data. Our implementation is highly modular and makes extensive use of unit testing to ensure the accuracy of results. This modularity allows alternative components for functionalities such as choosing operational taxonomic units (OTUs), sequence alignment, inferring phylogenetic trees and phylogenetic and taxon-based analysis of diversity within and between samples (including incorporation of third-party applications for many steps) to be easily integrated and benchmarked against one another (Supplementary Fig. 1).

We applied the QIIME workflow to a combined analysis of previously collected data (see Supplementary Discussion) for distal gut bacterial communities from conventionally raised mice, adult human monozygotic and dizygotic twins and their mothers, and a time series study of adult germ-free mice after they received human fecal microbiota (Fig. 1, Supplementary Table 1 and Supplementary Discussion). This analysis combined ten full 454 FLX runs and one partial run, totalling 3.8 million bacterial 16S rRNA sequences from previously published studies, including reads from different regions of the 16S rRNA gene.

QIIME is thus a robust platform for combining heterogeneous experimental datasets and for rapidly obtaining new insights about various microbial communities. Because QIIME scales to millions of sequences and can be used on platforms from laptops to high-performance computing clusters, we expect it to keep pace with advances in sequencing technology and to facilitate characterization of microbial community patterns ranging from normal variations to pathological disturbances in many human, animal and other environmental ecosystems.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank our collaborators for their helpful suggestions on features, documentation and the manuscript, and our funding agencies for their commitment to open-source software. This work was supported in part by Howard Hughes Medical Institute and grants from the Crohn’s and Colitis Foundation of America, the German Academic Exchange Service, the Bill and Melinda Gates Foundation, the Colorado Center for Biofuels and Biorefining and the US National Institutes of Health (DK78669, GM65103, GM8759, HG4872 and its ARRA supplement, HG4866, DK83981 and LM9451).

References

1. National Institutes of Health Human Microbiome Project Working Group *et al.* Genome Res. 2009; 19:2317–2323. [PubMed: 19819907]
2. Hopkin M. Nature. 2006; 444:420–421. [PubMed: 17122828]

3. Hamady M, Walker JJ, Harris JK, Gold NJ, Knight R. *Nat Methods*. 2008; 5:235–237. [PubMed: 18264105]
4. Cole JR, et al. *Nucleic Acids Res*. 2009; 37:D141–D145. [PubMed: 19004872]
5. Schloss PD, et al. *Appl Environ Microbiol*. 2009; 75:7537–7541. [PubMed: 19801464]
6. Knight R, et al. *Genome Biol*. 2007; 8:R171. [PubMed: 17708774]




Figure 1. QIIME analyses of the distal gut microbiotas of conventionally raised and conventionalized mice, gnotobiotic mice colonized with a human fecal gut microbiota (H-mice), and human adult mono- and dizygotic twins. (a) Principal coordinates analysis plots for mice, H-mice and twins. Colors correspond to separate samples by species and time point, and are consistent throughout the panels. (b) Unweighted UniFrac distance histograms between the data for fecal microbiota of human twins; human donors for the H-mice study; day 56 post-transplant H-mice on a low-fat (LF) and plant polysaccharide-rich (PP) diet; day 1 H-mice (LF and PP diet); and day 0 H-mice. Taxonomic classifications are presented at the class level. (c) Alpha diversity rarefaction plots of phylogenetic diversity for the H-mice samples. (d) OTU network connectivity of H-mice time series data. CONV-D, conventionalized mice; CONV-R, conventionally raised mice; and GF, germ-free mice.